




Contents lists available at ScienceDirect

Livestock Science

journal homepage: www.elsevier.com/locate/livsci

Effect of injectable copper, selenium, zinc and manganese on the pregnancy rate of crossbred heifers (*Bos indicus* × *Bos taurus*) synchronized for timed embryo transfer

J.N.S. Sales^a, R.V.V. Pereira^b, R.C. Bicalho^b, P.S. Baruselli^{a,*}

^a Department of Animal Reproduction, FMVZ-USP, Prof. Orlando Marques de Paiva, 87, CEP 05508-270, Sao Paulo, SP, Brazil

^b Department of Population Medicine and Diagnostic Sciences, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853, United States

ARTICLE INFO

Article history:

Received 9 December 2010

Received in revised form 12 April 2011

Accepted 21 June 2011

Available online xxxx

Keywords:

Cattle

Embryo transfer

Estrus synchronization

Trace mineral

ABSTRACT

The success of synchronization protocols for timed embryo transfer (TET) depends on various factors such as breed, body condition scores and nutrition (including trace mineral status). In the present study, crossbred heifers were treated with subcutaneous injectable trace minerals (ITM – 100 mg zinc, 100 mg manganese, 50 mg copper and 25 mg selenium; Multimin®, Minerthal, Brazil) 17 days prior to embryo transfer. Estrus synchronization for TET, conception rate and pregnancy loss were evaluated. Subcutaneous administration of ITM did not increase the number of heifers successfully synchronized compared to non-injected controls [ITM group = 82.1% (308/375) versus control = 83.1% (375/451); $P=0.76$]. However, heifers belonging to the ITM group ($n=219$) had a 1.58 fold and 1.72-fold higher ($P=0.005$) odds of being pregnant 23 and 48 days after TET compared to the control group ($n=276$; not receiving ITM), respectively. In contrast, treatment with subcutaneous ITM did not significantly lower the pregnancy-loss rate [control group = 17.2% (17/99) versus ITM group = 10.6% (11/105); $P=0.18$]. Even though subcutaneous administration of ITM 17 days prior to TET did not increase the number of heifers successfully synchronized, a significant increase in the conception rate (embryo survival) at 23 and 48 days after TET was noted in heifers of the ITM group.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Timed embryo transfer (TET) protocols are well established in the dairy and beef industry. Several studies have been performed in order to reduce management problems that lessen the efficiency of TET, thus making the procedure financially viable (Baruselli et al., 2010; Binelli et al., 2001; Block et al., 2003; Bo et al., 2002). However, less effort has been directed toward investigating the effects of nutrition on TET, even though studies have clearly established a profound influence of nutrition on the reproductive performance of female cattle (Campanile et al., 2010; Robinson, 1996; Sales et al., 2008; Webb et al., 1999). Moreover, the effects of minerals in improving fertility have also been described (Wilde,

2006). Ballantine et al. (2002) observed that supplementation with Cu, Zn, Mn and Co (inorganic versus organic) reduced days open and tended to improve both the first-service conception rate and the percentage of cows pregnant at 150 days postpartum. Furthermore, the concept that certain essential trace minerals play crucial roles in enzymatic and metabolic pathways which are critical for embryo development during pregnancy in cattle has been described (Griffiths et al., 2007; Hostetler et al., 2003). The embryo relies entirely on the maternal system for its supply of trace minerals and other nutrients needed for normal development. Hostetler et al. (2003) observed that the levels of zinc, copper and manganese in the conceptus were severalfold greater than in the surrounding reproductive tissues, indicating preferential accumulation of these minerals by the conceptus. Trace minerals can be supplied to cattle by various means, including as free-choice minerals, drenches and pastes (Ahola et al., 2004). The

* Corresponding author. Tel./fax: +55 11 3091 7674.

E-mail address: barusell@usp.br (P.S. Baruselli).

recent development of injectable chelated minerals introduces a new option for supplementing these essential minerals.

The objective of this study was to evaluate the effect of injectable trace mineral (ITM) supplementation (100 mg zinc, 100 mg manganese, 50 mg copper and 25 mg selenium) on the number of heifers successfully synchronized, and on the conception and pregnancy-loss rates of embryo recipients submitted to TET protocols. The hypothesis of this study was that ITM supplementation increases both the number of heifers successfully synchronized and the conception rate, and reduces the pregnancy-loss rate of crossbred heifers submitted to TET protocols.

2. Materials and methods

2.1. Study animals, facilities and diets

The study was performed from December 2007 to November 2009 at five different farms in Brazil. At each farm the heifers received free-choice oral mineral supplementation offered according to nutritional requirements of the specific animal category (NRC, 2001). In total, 826 crossbred heifers (*Bos indicus* × *Bos taurus*) cycling (presence of the corpus luteum; CL), presenting a body condition score (BCS) of 2.75 to 3.5 (1 to 5 scale, Wildman et al., 1982) and a body weight of 300–320 kg were enrolled in the study. During the summer, the animals were maintained in *Brachiaria brizantha* grass paddocks in a rotational grazing system. In winter, the heifers were supplemented with sugarcane and a protein and mineral supplement to compensate for the low nutritional value of the forage. During both seasons (summer and winter), the animals had ad libitum access to water and mineral salts (Minerthal 160 Corte®, Minerthal, Brazil; Table 1). The protein and mineral supplement was composed of 10% mineral salts, 10% urea, 17.5% salt, 47.5% cotton seed and 15% citrus pulp and was provided every 2 days. The expected daily intake was 75 g of mineral salt (summer) or 400 g of protein and mineral supplement (winter). All minerals were supplied at NRC recommended levels (NRC, 2001).

2.2. Estrus synchronization for timed embryo transfer

At unknown stages of the estrous cycle (D0), all animals received 2 mg of estradiol benzoate (Sincrodiol®, Ouro Fino,

Brazil) and one progesterone-releasing intravaginal device (Sincrogest®, Ouro Fino, Brazil). Eight days later (D8), the intravaginal device was removed and 500 µg of cloprostenol (Sincrocio®, Ouro Fino, Brazil), 300 IU of equine chorionic gonadotropin (eCG) (Novormon®, Syntex, Argentina) and 1.0 mg of estradiol cypionate (SincroCP®, Ouro Fino, Brazil) were injected in all animals. Embryo transfer was performed on day 17. Only grade 1 in vitro-derived *B. indicus* embryos (Stringfellow and Seidel, 1998) were used. Pregnancy diagnosis was performed using ultrasonography at 23 and 48 days after TET. An ultrasound machine with an 8 MHz crystal array vet linear transducer Aquila Probe (Pie Medical, ESAOTE, The Netherlands) was used.

2.3. Treatment groups

The heifers were randomly allocated to one of two treatments with an allocation ratio of 1:1 in a blinded fashion. The randomization was stratified by farm and BCS. The experimental groups were: control (n = 276), where animals did not receive any kind of ITM supplementation, and ITM (n = 219), where animals received 5 ml subcutaneous injection of copper, manganese, zinc and selenium (Multimin®, Minerthal, Brazil). The injection was administered at the onset of the TET synchronization protocol (D0).

This trial was conducted in agreement with the ethical principles in animal research adopted by the Bioethics Commission of the School of Veterinary Medicine and Animal Science of the University of São Paulo, Brazil.

2.4. Statistical analysis

Synchronization and pregnancy rates were summarized in tabular format using frequencies and percentages. The chi-square test was used to compare synchronization rates between groups. Pregnancy rates were analyzed using the intent-to-treat population, which was defined as all animals which were successfully synchronized and received an embryo. Logistic regression analysis was performed to compare all study outcomes (pregnant at day 23 after TET, pregnant at day 48 after TET, pregnancy loss) between the two study arms. In order to adjust the analysis for farm site and year effect, those variables were included as covariates in the logistic regression analysis models. All *P*-values are two-sided, with *P* < 0.05 indicating a statistically significant difference. Data analysis was performed using SAS® version 9.2 software (SAS Corp. Cary, NC).

3. Results

Of the 826 heifers enrolled for synchronization, only 495 (control n = 276 and ITM n = 219) received an embryo. Despite the fact that approximately 80% of heifers had been successfully synchronized (presence of CL on D17), the number of embryos produced in vitro was insufficient to allow enough embryos to be transferred to all synchronized animals.

There was no interaction (*P* = 0.94) between the treatment, BCS, year and farm. Furthermore, no statistical difference between the number of heifers successfully synchronized and ready for TET was observed between animals in the control

Table 1

Mineral salt composition. Levels of guarantee per kg of product.

Element	Mineral salt
Phosphorus (g)	90
Calcium (g)	160
Magnesium (g)	15
Sulfur (g)	15
Sodium (g)	141
Zinc (mg)	3600
Copper (mg)	960
Manganese (mg)	1020
Cobalt (mg)	100
Iodine (mg)	63
Selenium (mg)	13
Fluorine (mg)	900

Table 2

Outcome of logistic regression evaluating the effect of treatment group and farm effect on the odds of conception and on conception rate at 23 and 48 days post embryo transfer.

Variables	CR* at 23 days	OR**	P	CR* at 48 days	OR**	P
<i>Group</i>						
Control ^a	36% (276)	Ref	0.015	30% (276)	Ref	0.005
ITM ^b	48% (219)	1.58		43% (219)	1.72	
<i>Farm</i>						
Farm A	32% (240)	Ref	0.0007	28% (240)	Ref	0.007
Farm B	37% (27)	1.18		33% (27)	1.16	
Farm C	45% (53)	1.74		34% (53)	1.25	
Farm D	53% (78)	2.30		51% (78)	2.56	
Farm E	55% (97)	2.55		42% (97)	1.80	

*CR = conception rate at 23 or 48 days of gestation; **OR = odds ratio.

^a Control group not receiving injection of trace minerals.

^b Treatment group receiving an injection of trace minerals 17 days prior to embryo transfer.

[83.1% (375/451)] or ITM group [82.1% (308/375); $P=0.76$]. Comparison of conception rates between the ITM and control groups showed significant differences both 23 days after TET (48% vs. 36%, $P=0.015$) and 48 days after TET (43% vs. 30%, $P=0.005$) (Table 2). There was a significant farm effect (Table 2) but no significant year effect. The conception rate for cows diagnosed pregnant 23 days after TET across the five farms ranged from 32% to 55% ($P=0.0007$) and for cows diagnosed pregnant 48 days after TET ranged from 28% to 51% ($P=0.007$). A logistic regression evaluating the effect of treatment group for pregnancy loss showed that the control group had a pregnancy loss of 17% compared to 11% for the ITM group (Table 3).

4. Discussion

The hypothesis that ITM supplementation would increase the number of synchronized heifers submitted to TET protocols was not confirmed. Although no difference was noted in the rate of successful synchronization, significant differences were observed in pregnancy rates both at 23 days after TET and at

Table 3

Outcome of logistic regression evaluating the effect of treatment group (control or ITM) and farm on the odds of pregnancy loss.

Variables	PL*	OR**	P
<i>Group</i>			
Control ^a	17.2% (99)	1.78	0.18
ITM ^b	10.6% (105)	Ref	
<i>Farm</i>			
Farm A	11% (76)	Ref	0.06
Farm B	10% (10)	1.00	
Farm C	25% (24)	2.87	
Farm D	2% (41)	0.22	
Farm E	23% (53)	2.53	

*PL – Pregnancy loss that occurred between the first and second pregnancy checks; **OR – odds ratio.

^a Control group not receiving injection of trace minerals.

^b Treatment group receiving an injection of trace minerals 17 days prior to embryo transfer.

48 days after TET, with the ITM group presenting 1.58-fold and 1.72-fold higher odds of conception rate at 23 and 48 days when compared to the control group (Table 2; $P=0.005$), respectively. These findings are consistent with related reproductive improvements observed in other studies, in which mineral supplementation improved cow fertility (Ahola et al., 2004; Moeini et al., 2009). In addition, Ballantine et al. (2002) observed that supplementation with Cu, Zn, Mn and Co reduced days open without changing the first-service conception rate. In contrast, Lamb et al. (2008) reported that mineral supplementation did not alter embryo number or quality in superovulated Angus heifers.

While fertilization rates in cows can be as high as 80%, the rate of embryonic mortality during the first 30 days of pregnancy is on average about 40% (Santos et al., 2004). A review found that much of the pregnancy loss is concentrated in the embryonic period, during the first 42 days after breeding (Inskeep and Dailey, 2005).

The positive effects of mineral supplementation found in the present study could perhaps be associated with mineral action on the immune response and cellular enzymatic activities, including hormone production (Bendich, 1993; Hostetler et al., 2003; Niederman et al., 1994; Salman et al., 2009). Although there is known deficiency of P, Co, Cu, I, Zn and Se in the geographic areas of the study (Tokarnia et al., 1999), the animals that received mineral supplementation likely acquired sufficient amounts of trace minerals for synthesis of proper levels of hormones. This supplementation could have contributed to the higher conception rate in heifers submitted to TET in the ITM group. Retention of pregnancy in the cow relies on the maintenance of the corpus luteum and consequent progesterone secretion, one of the main hormones responsible for the maintenance of pregnancy (Binelli et al., 2001). Estradiol also has been demonstrated to have a positive effect on pregnancy retention (Starbuck et al., 2004). Moreover, copper is an integral part of superoxide dismutase (SOD) enzyme systems, and an interaction has been shown between zinc deficiency and a depressed immune system (Engle et al., 1997; Enjalbert et al., 2006; Nocek et al., 2006; Prasad and Kundu, 1995). Manganese is another important trace mineral whose deficiency can interfere with reproduction. Manganese acts as a cofactor in the synthesis of cholesterol, a precursor of steroids, including estrogen and progesterone (Nocek et al., 2006). Furthermore, manganese is a structural component of manganese superoxide dismutase (MnSOD), which regulates cellular differentiation during early embryonic development (Hostetler et al., 2003). Selenium, another trace mineral, is an essential element of several enzyme systems, including glutathione-peroxidase (GSHpx) and deiodinase (Burk and Hill, 1993; Spears and Weiss, 2008). Both these enzymes have been shown to be important in embryonic development, through their antioxidant functions and regulatory effects on metabolic rate (Hostetler et al., 2003). Moreover, selenium is important in inhibiting nitric oxide production in cells (an important controller of steroidogenesis in and apoptosis of granulosa cells), in stimulating proliferation of the cells of small follicles, and in increasing estrogen output by follicles, all of which could act to augment pregnancy retention in the cow (Basini and Tamanini, 2000; Faes et al., 2009; Starbuck et al., 2004).

Zinc is a cofactor in DNA synthesis and gene transcription (Chesters, 1978). It also is essential for the activity of copper/zinc SOD and has a modulating action on IGF at the cellular level (Zapf and Froesch, 1986). Both these enzymes are present at high levels in the uterus during early pregnancy (Hostetler et al., 2000). The fact that zinc, copper and manganese are present at severalfold higher levels in the conceptus compared to other reproductive tissues indicates the importance of supplying these micronutrients in sufficient amounts to the cow (Hostetler et al., 2003). Based on this study it can be inferred that supplementation of copper, zinc, manganese and selenium via the injectable route could increase pregnancy rates by increasing early embryonic survival.

In summary, subcutaneous administration of ITM 17 days prior to TET did not increase the number of heifers successfully synchronized, although it did increase the conception rates (embryo survival) 23 and 48 days after TET in crossbred heifers submitted to TET.

Acknowledgments

We thank Professor Jens Eickhoff from Colorado State University in Fort Collins for his diligent analysis and reporting on the statistical section of the manuscript and Paraíso Embrões and Ventre Vivo companies. This research was supported by Multimin Co.

References

- Ahola, J.K., Baker, D.S., Burns, P.D., Mortimer, R.G., Enns, R.M., Whittier, J.C., Geary, T.W., Engle, T.E., 2004. Effect of copper, zinc, and manganese supplementation and source on reproduction, mineral status, and performance in grazing beef cattle over a two-year period. *J. Anim. Sci.* 82, 2375–2383.
- Ballantine, H.T., Socha, M.T., Tomlinson, D.J., Johnson, A.B., Fielding, A.S., Shearer, J.K., van Amstel, S.R., 2002. Effects of feeding complexed zinc, manganese, copper and cobalt to late gestation and lactating dairy cows on claw integrity, reproduction and lactation performance. *Prof. Anim. Sci.* 18, 211–218.
- Baruselli, P.S., Ferreira, R.M., Sá Filho, M.F., Nasser, L.F.T., Rodrigues, C.A., Bó, G.A., 2010. Bovine embryo transfer recipient synchronization and management in tropical environments. *Reprod. Fert. Dev.* 22, 67–74.
- Basini, G., Tamanini, C., 2000. Selenium stimulates estradiol production in bovine granulosa cells: possible involvement of nitric oxide. *Domest. Anim. Endocrinol.* 18, 1–17.
- Bendich, A., 1993. Physiological role of antioxidants in the immune system. *J. Dairy Sci.* 76, 2789–2794.
- Binelli, M., Thatcher, W.W., Mattos, R., Baruselli, P.S., 2001. Antiluteolytic strategies to improve fertility in cattle. *Theriogenology* 56, 1451–1463.
- Block, J., Drost, M., Monson, R.L., Rutledge, J.J., Rivera, R.M., Paula-Lopes, F.F., Ocon, O.M., Krininger, C.E., Liu, J., Hansen, P.J., 2003. Use of insulin-like growth factor-I during embryo culture and treatment of recipients with gonadotropin-releasing hormone to increase pregnancy rates following the transfer of in vitro-produced embryos to heat-stressed, lactating cows. *J. Anim. Sci.* 81, 1590–1602.
- Bó, G.A., Baruselli, P.S., Moreno, D., Cutaia, L., Caccia, M., Tribulo, R., Tribulo, H., Mapletoft, R.J., 2002. The control of follicular wave development for self-appointed embryo transfer programs in cattle. *Theriogenology* 57, 53–72.
- Burk, R.F., Hill, K.E., 1993. Regulation of selenoproteins. *Annu. Rev. Nutr.* 13, 65–81.
- Campanile, G., Baruselli, P.S., Vecchio, D., Prandic, A., Neglia, G., Carvalho, N.A.T., Sales, J.N.S., Gasparinina, B., D'Occhio, M.J., 2010. Growth, metabolic status and ovarian function in buffalo (*Bubalus bubalis*) heifers fed a low energy or high energy diet. *Anim. Reprod. Sci.* 122, 74–81.
- Chesters, J.K., 1978. Biochemical functions of Zn in animals. *World Rev. Nutr. Diet.* 32, 135–164.
- Engle, T.E., Nockels, C.F., Kimberling, C.V., Weaber, D.L., Johnson, A.B., 1997. Zinc repletion with organic or inorganic forms of zinc and protein turnover in marginally zinc-deficient calves. *J. Anim. Sci.* 75, 3074–3081.
- Enjalbert, F., Lebreton, P., Salat, O., 2006. Effects of copper, zinc and selenium status on performance and health in commercial dairy and beef herds: retrospective study. *J. Anim. Physiol. Anim. Nutr.* 90, 459–466.
- Faes, M.R., Caldas-Bussiére, M.C., Viana, K.S., Dias, B.L., Costa, F.R., Escocard, R.M., 2009. Nitric oxide regulates steroid synthesis by bovine antral granulosa cells in a chemically defined medium. *Anim. Reprod. Sci.* 110, 222–236.
- Griffiths, L.M., Loeffler, S.H., Socha, M.T., Tomlinson, D.J., Johnson, A.B., 2007. Effects of supplementing complexed zinc, manganese, copper and cobalt on lactation and reproductive performance of intensively grazed lactating dairy cattle on the South Island of New Zealand. *Anim. Feed Sci. Technol.* 137, 69–83.
- Hostetler, C.E., Cronrath, J.D., Becker, W.C., Miranda, M.A., 2000. Dietary supplementation of proteinated trace minerals (OPTiMIN) in sows and replacement gilts increases mineral concentrations in reproductive tissues. 14th International Congress Animal Reproduction 1, 272, Abstr.
- Hostetler, C.E., Kincaid, R.L., Miranda, M.A., 2003. The role of essential trace elements in embryonic and fetal development in livestock. *Vet. J.* 166, 125–139.
- Inskeep, E.K., Dailey, R.A., 2005. Embryonic death in cattle. *Vet. Clin. North Am. Food Anim. Pract.* 21, 437–461.
- Lamb, G.C., Brown, D.R., Larson, J.E., Dahlen, C.R., DiLorenzo, N., Arthington, J.D., DiCostanzo, A., 2008. Effect of organic or inorganic trace mineral supplementation on follicular response, ovulation, and embryo production in superovulated Angus heifers. *Anim. Reprod. Sci.* 106, 221–231.
- Moeini, M.M., Karami, H., Mikaeli, E., 2009. Effect of selenium and vitamin E supplementation during the late pregnancy on reproductive indices and milk production in heifers. *Anim. Reprod. Sci.* 114, 109–114.
- Niederman, C.N., Blodgett, D., Eversole, D., Schurig, G.G., Thatcher, C.D., 1994. Effect of copper and iron on neutrophil function and humoral immunity of gestating beef cattle. *J. Am. Vet. Med. Assoc.* 204, 1796–1800.
- Nocek, J.E., Socha, M.T., Tomlinson, D.J., 2006. The effect of trace mineral fortification level and source on performance of dairy cattle. *J. Dairy Sci.* 89, 2679–2693.
- Prasad, T., Kundu, M.S., 1995. Serum IgG and IgM responses to sheep red blood cells (SRBC) in weaned calves fed milk supplemented with Zn and Cu. *Nutrition* 11, 712–715.
- Robinson, J.J., 1996. Nutrition and reproduction. *Anim. Reprod. Sci.* 42, 25–34.
- Sales, J.N.S., Dias, L.M.K., Viveiros, A.T.M., Pereira, M.N., Souza, J.C., 2008. Embryo production and quality of Holstein heifers and cows supplemented with betacarotene and tocopherol. *Anim. Reprod. Sci.* 106, 77–89.
- Salman, S., Khol-Parisini, A., Schaff, H., Lahrssen-Wiederholt, M., Hulan, H.W., Dinse, D., Zentek, J., 2009. The role of dietary selenium in bovine mammary gland health and immune function. *Anim. Health Res. Rev.* 10, 21–34.
- Santos, J.E., Thatcher, W.W., Chebel, R.C., Cerri, R.L., Galvao, K.N., 2004. The effect of embryonic death rates in cattle on the efficacy of estrus synchronization programs. *Anim. Reprod. Sci.* 82–83, 513–535.
- Spears, J.W., Weiss, W.P., 2008. Role of antioxidants and trace elements in health and immunity of transition dairy cows. *Vet. J.* 176, 70–76.
- Starbuck, M.J., Dailey, R.A., Inskeep, E.K., 2004. Factors affecting retention of early pregnancy in dairy cattle. *Anim. Reprod. Sci.* 84, 27–39.
- Stringfellow, D.A., Seidel, S.M., 1998. Manual of the International Embryo Transfer Society: a Procedural Guide and General Information for the Use of Embryo Transfer Technology, Emphasizing Sanitary Procedures, third ed. Savory III, USA.
- Tokarnia, C.H., Jürgen, D.M., Sheila, S., Peixoto, P.V., 1999. Mineral deficiencies and imbalances in cattle and sheep: a review of Brazilian studies made between 1987 and 1998. *Braz. J. Vet. Res.* 19, 47–62.
- Webb, R., Royal, M.D., Gong, J.G., Garnsworthy, P.C., 1999. The influence of nutrition on fertility. *Pract. Pract.* 7, 227–234.
- Wilde, D., 2006. Influence of macro and micro minerals in the peri-parturient period on fertility in dairy cattle. *Anim. Reprod. Sci.* 96, 240–249.
- Wildman, E.E., Jones, G.M., Wagner, P.E., Boman, R.L., Troutt Jr., H.F., Leschb, T.N., 1982. A dairy cow body condition scoring system and its relationship to selected production characteristics. *J. Dairy Sci.* 65, 495–501.
- Zapf, J., Froesch, R., 1986. Insulin-like growth factors/somatomedins: structure, secretion, biological actions and physiological role. *Horm. Res.* 24, 121–130.